

## §6-2—6-3 Volumes by Cylindrical Shells


\*公式：

$$V_3 = \int V_2 dx$$

$$V_3 = \begin{cases} \text{● ○ §6-2(截面為圓環)} \\ \text{○ §6-3(截面為圓柱的表面)} \end{cases}$$

**Example 1** : Prove that the volume of the sphere with radius  $r$  is  $\frac{4}{3}\pi r^3$ .


**Solution** :



$$V = 2 \int_0^r \pi r'^2 dx = 2 \int_0^r \pi (r^2 - x^2) dx$$

$$= \frac{4}{3} \pi r^3$$

§6-2



$$V = 2 \int_0^r 2\pi \times x h dx$$


$$= 2 \int_0^r 2\pi \times x \sqrt{r^2 - x^2} dx$$

$$= -\frac{4\pi}{3} (r^2 - x^2)^{\frac{3}{2}} \Big|_0^r$$

$$= \frac{4}{3} \pi r^3$$

§6-3

**Example 2 :** Find the volume of the following graph.



**Solution :**


$$\frac{x}{h} = \frac{r'}{r}$$

$$\Rightarrow r' = \frac{r}{h}x.$$


$$V = \int_0^h \pi r'^2 dx = \int_0^h \pi \left( \frac{r^2}{h^2} \right) x^2 dx$$

$$= \frac{1}{3} \pi r^2 h.$$

**Example 3 :** A frustum of a pyramid with square base of side  $b$ , square top of side  $a$ , and height  $h$ .



**Solution :**



$$V = \int_0^h r^2 dx$$

$$\frac{k}{h} = \frac{a}{b-a} \Rightarrow k = \frac{ah}{b-a}$$


$$\frac{x+k}{k} = \frac{r}{a} \Rightarrow r = a + \left(\frac{b-a}{h}\right)x$$

$$\Rightarrow V = \int_0^h \left(a + \frac{b-a}{h}x\right)^2 dx$$


$$= \frac{h}{3(b-a)} \left(a + \frac{b-a}{h}x\right)^3 \Big|_0^h$$

$$= \frac{h}{3}(a^2 + ab + b^2).$$

**Example 4 :** Rotating  $\begin{cases} y = x \\ y = x^2 \end{cases}$  about the  $x$ -axis.



**Solution :**



Method 1 : (截面為環狀)

$$V = \int_0^1 (\pi r_1^2 - \pi r_2^2) dx, \quad r_1 = x, r_2 = x^2.$$

$$= \int_0^1 (\pi x^2 - \pi x^4) dx = \frac{2\pi}{15}$$


Method 2 : (截面為圓柱狀)

$$V = \int_0^1 2\pi r h dy$$

$$= \int_0^1 2\pi y (\sqrt{y} - y) dy.$$

**Example 5 :** Rotating  $\begin{cases} y = 8 \\ y = x^3 \\ x = 0 \end{cases}$  about the  $x$ -axis.

**Solution :**




$$V = \int_0^2 (64 - x^6) \pi dx \text{ (截面為環狀)}$$

$$= \int_0^8 2\pi y^{\frac{1}{3}} y dy \text{ (截面為圓柱狀)}$$

$$= \frac{768}{7} \pi.$$

**Example 6 :** Rotating  $\begin{cases} y = x - x^2 \\ y = 0 \end{cases}$  about the line  $x = 2$ .


**Solution :**



$$V = \int_0^1 2\pi(2-x)(x-x^2) dx$$




$$= \frac{\pi}{2}.$$

**Example 7 :** A wedge is cut out of a circular cylinder of radius 4 by two planes. One plane is perpendicular to the axis of the cylinder. The other intersects the first at an angle of  $30^\circ$  along a diameter of the cylinder. Find the volume of the wedge.



**Solution :**

Method 1 :






$$V = \int_0^4 \text{長方形截面} dx$$

$$= \int_0^4 \underbrace{2\sqrt{4^2 - x^2}}_{\text{長}} \underbrace{\left(\frac{x}{\sqrt{3}}\right)}_{\text{寬}} dx$$

$$= \frac{128}{3\sqrt{3}}$$

Method 2 :



$$V = \int_0^4 \Delta ABC \text{截面} dx$$


$$= \int_0^4 \overline{AB} \times \overline{BC} dx$$

$$= \int_0^4 \frac{16 - x^2}{\sqrt{3}} dx$$

$$= \frac{128}{3\sqrt{3}}$$

**Example 8 :** Find the volume common to two spheres, each with radius  $r$ . If the center of each sphere lies on the surface of other.

**Solution :**



**Method 1 :**

$$\begin{aligned} V &= 2 \int_0^r \pi y^2 dx \\ &= 2 \int_0^r \pi \left( r^2 - \left( x + \frac{r}{2} \right)^2 \right) dx \\ &= \frac{5}{24} \pi r^3. \end{aligned}$$

**Method 2 :**


$$\begin{aligned} V &= 2 \int_{-\frac{r}{2}}^0 \pi y^2 dx \\ &= 2 \int_{-\frac{r}{2}}^0 \pi \left( r^2 - \left( x - \frac{r}{2} \right)^2 \right) dx \\ &= \frac{5}{24} \pi r^3. \end{aligned}$$

**Example 9 :** Find the volume common to two circular cylinders, each with radius  $r$ . If the axes of the cylinders intersect at right angle.

\*想一想：

1. 先想想看，交集區域或是交集區域集面樣子？
2. 若想不出來，試著用代數手法組成此交集區域。

**Solution :**



由對稱性，我們只考慮交集的 $\frac{1}{8}$ 區域，即  $x \geq 0, y \geq 0, z \geq 0$ .

- 當  $z = 0 \Rightarrow 0 \leq x \leq r, 0 \leq y \leq r$

(正方形，邊長 =  $r$ )

- 當  $z = r \Rightarrow x = 0, y = 0$

(正方形，邊長 = 0)

- 當  $0 < z < r \Rightarrow 0 \leq x \leq \sqrt{r^2 - z^2}, 0 \leq y \leq \sqrt{r^2 - z^2}$

(正方形，邊長 =  $\sqrt{r^2 - z^2}$ )

$\Rightarrow$  當  $z$  固定， $0 \leq z \leq r$ ，交集區域的橫截面為正方形。

$$\Rightarrow \text{交集區域的體積 } V = 8 \int_0^r (r^2 - z^2) dz = \frac{16}{3} r^3.$$